

## HÄMI Referensdatabas – Biologiska mätdata

*Uppdaterad 2010-11-02*

Aberg A, Tysklind M, Nilsson T, Macleod M, Hanberg A, Andersson R, Bergek S, Lindberg R, Wiberg K. [Exposure assessment at a PCDD/F contaminated site in Sweden-field measurements of exposure media and blood serum analysis](#). Environ Sci Pollut Res Int. 2010 Jan;17(1):26-39. Epub 2009 Jul 30.

Alfvén T, Elinder CG, Carlsson MD, Grubb A, Hellström L, Persson B, Pettersson C, Spang G, Schütz A, Järup L (2000) Low-level cadmium exposure and osteoporosis. Bone Miner Res 15:1579-86

Alfvén T (2002) Bone and kidney effects from cadmium exposure. Akademisk avhandling. Karolinska Institutet

Alfven T, Elinder CG, Hellström L, Lagarde F, Järup L. Cadmium exposure and distal forearm fractures. J Bone Miner Res 2004;19(6):900–5.

Andersson C (1998) Kvicksilverbelastning hos Hagforsbor. Examensarbete, 20 poäng. Miljö- och hälsoskyddsprogrammet, Umeå Universitet

Andersson C, Sjörs A (1998) Kvicksilverbelastning hos Hagforsbor. Kvicksilver i Värmländska sjöar. Hagfors kommun

Andersson C, Sällsten G, Schütz A, Barregård L (2000) Samband mellan konsumtion av insjöfisk och kvicksilverbelastning hos personer utan amalgamfyllning. Rapport från Yrkes- och miljömedicin, Göteborg, nr 78

Andersson K, Andersson I, Botvalde M, Hjelm L, Fredriksson R, Hogstedt C, Oliv Å, Sandegren G, Sundell L, Ulander A (1981) Studie av referensvärden för bly i blod och urin hos vuxna. Läkartidningen. 78. 39, 3378-3379

Andrén P, Schütz A, Vahter M, Attewell R, Johansson L, Willers S, Skerfving S (1988) Environmental exposure to lead and arsenic among children living near a glassworks. Sci Tot Environ 77:25-34

Ask K, Åkesson A, Berglund M, Vahter M (2002) Inorganic mercury and methylmercury in placentas of Swedish women. Environ Health Perspect 110:523-6

Asplund L, Svensson BG, Nilsson A, et al (1994) Polychlorinated biphenyls, 1,1,1-trichloro-2,2-bis(p-chlorophenyl)ethane (p,p'-DDT) and 1,1-dichloro-2,2-bis(p-chlorophenyl)-ethylene (p,p'-DDE) in human plasma related to fish consumption. Arch Environ Health 49:477-486

Attewell R, Schütz A, Skerfving S, Andrén P, Willers S (1988) Samhällets åtgärder mot blyexponering har sänkt blodblyhalterna markant. *Läkartidningen*. 85. 30-31, 2458-2462

Axmon A, Rylander L, Strömberg U, Dyremark E, Hagmar L (2001) Polychlorinated biphenyls in blood plasma among Swedish female fish consumers in relation to time to pregnancy. *J Toxicol Environ Health A* 23;64:485-98

Axmon A, Rylander L, Strömberg U, Hagmar L (2000) Miscarriages and stillbirths in women with a high intake of fish contaminated with persistent organochlorine compounds. *Int Arch Occup Environ Health* 73:204-8

Axmon A, Rylander L, Strömberg U, Jönsson B, Nilsson-Ehle P, Hagmar L (2004) Polychlorinated biphenyls in serum and time to pregnancy. *Environmental Research* 96:186-195

Axmon A, Hagmar L, Jönsson BA. Rapid decline of persistent organochlorine pollutants in serum among young Swedish males. *Chemosphere* 2008 Feb;70(9):1620-8.

Baecklund M (1996) Environmental and genetic factors determining the concentration of cadmium and lead in blood from middle aged and elderly people. Examensarbete, 20 poäng. IMM, Karolinska Institutet

Baecklund M, Pedersen NL, Björkman L, Vahter M (1999) Variation in blood concentrations of cadmium and lead in the elderly. *Environ Res* 80:222-30

Barany E (2002) Trace elements in adolescents. Akademisk avhandling, Sveriges Lantbruksuniversitet

Barany E, Bergdahl IA, Bratteby LE, Lundh T, Samuelson G, Schütz A, Skerfving S, Oskarsson A (2002) Relationships between trace element concentrations in human blood and serum. *Toxicol Lett* 5:177-84

Barany E, Bergdahl IA, Bratteby LE, Lundh T, Samuelson G, Schütz A, Skerfving S, Oskarsson A (2002) Trace element levels in whole blood and serum from Swedish adolescents. *Sci Total Environ* 8;286:129-41

Barany E, Bergdahl IA, Bratteby LE, Lundh T, Samuelson G, Schütz A, Skerfving S, Oskarsson A (2002) Trace elements in blood and serum of Swedish adolescents: relation to gender, age, residential area, and socioeconomic status. *Environ Res* 89:72-84

Barany E, Bergdahl IA, Bratteby LE, Lundh T, Samuelson G, Skerfving S, Oskarsson A (2003) Mercury and selenium in whole blood and serum in relation to fish consumption and amalgam fillings in adolescents. *J Trace Elem Med Biol* 17:165-70

Barregård L, Svalander C, Schütz A, Westberg G, Sällsten G, Blohmé I, Mölne J, Attman P-O, Haglind P (1999) Cadmium, mercury and lead in kidney cortex of the general

Swedish population: a study of biopsies from living kidney donors. *Environ Health Perspect* 107:867-871

Barregård L, Sällsten G, Törnström G (1996) Kvicksilver i blod hos gravida kvinnor i Västsverige. Rapport från YMK i Göteborg nr 62

Barregard L, Fabricius-Lagging E, Lundh T, Mölne J, Wallin M, Olausson M, Modigh C, Sallsten G. [Cadmium, mercury, and lead in kidney cortex of living kidney donors: Impact of different exposure sources.](#) *Environ Res.* 2010 Jan;110(1):47-54.PMID: 19931045

Bensryd I, Rylander L, Högstedt B, Aprea P, Bratt I, Fahraeus C, Holmen A, Karlsson A, Nilsson A, Svensson BL, et al (1994) Effect of acid precipitation on retention and excretion of elements in man. *Science Tot Environ* 145:81-102

Bergdahl IA, Gerhardsson L, Schütz A, Desnick RJ, Wetmur JG, Skerfving S (1997) Delta-aminolevulinic acid dehydratase polymorphism: influence on lead levels and kidney function in humans. *Arch Environ Health* 52:91-6

Bergdahl IA, Skerfving S (1997) Partition of circulating lead between plasma and red cells does not seem to be different for internal and external sources of lead. *Am J Ind Med* 32:317-8

Bergdahl IA, Vahter M, Counter SA, Schütz A, Buchanan LH, Ortega F, Laurell G, Skerfving S (1999) Lead in plasma and whole blood from lead-exposed children. *Environ Res* 80:25-33

Berger U, Glynn A, Holmström KE, Berglund M, Ankarberg EH, Törnkvist A. [Fish consumption as a source of human exposure to perfluorinated alkyl substances in Sweden - analysis of edible fish from Lake Vättern and the Baltic Sea.](#) *Chemosphere.* 2009 Aug;76(6):799-804. Epub 2009 May 19.PMID: 19457539

Berglund M, Fahlgren L, Freland M, Vahter M (1994) Metaller i mark i Stockholms innerstad och kranskommuner –förekomst och hälsorisker för barn. IMM-rapport 2/94

Berglund M, Lind B, Björnberg KA, Palm B, Einarsson Ö, Vahter M (2005) Inter-individual variations of human mercury exposure biomarkers: a cross-sectional assessment. *Environ Health* 3;4:20

Berglund M, Lind B, Lannerö E, Vahter M (1994) A pilot study of lead and cadmium exposure in young children in Stockholm, Sweden: methodological considerations using capillary blood microsampling. *Arch Environ Cont Toxicol* 27:281-287

Berglund M, Lind B, Sörensen S, Vahter M (2000) Impact of soil and dust lead on children's blood lead in contaminated areas of Sweden. *Arch Environ Health* 55(2):93-7

Berglund M, Vahter M, Slorach S (1991) Determination of human exposure to lead and cadmium a WHO-UNEP pilot study. *Chem Spec Bioavail* 3: 69-72

Berglund M, Åkesson A, Lind B, Björs U, Palm B, Vahter M (1998) Förekomst av tungmetaller hos gravida kvinnor i Solna. I Naturvårdsverket rapport 4902

Berglund M, Åkesson A, Nermell B, Vahter M (1994) Intestinal absorption of dietary cadmium in women depends on body iron stores and fiber intake. *Environ Health Perspect* 102:1058-66

Bjerre B, Berglund M, Harsbo K, Hellman B (1993) Blood lead concentrations of Swedish preschool children in a community with high lead levels from mine waste in soil and dust. *Scand J Work Environ Health* 19:154-61

Bjerre B, Berglund M, Hellman B, Harsbo K (1998) Blodblyundersökning av förskolebarn - Inom område med höga metallhalter i Falu tätort. Miljö- och hälsoskydd i Falu kommun, MH 1998:6

Björkman L, Vahter M, Pedersen NL (2000) Both the environment and genes are important for concentrations of cadmium and lead in blood. *Environ Health Perspect* 108:719-22

Björnberg KA (2005) Mercury exposure during early human development. Akademisk avhandling. Karolinska Institutet

Björnberg KA, Vahter M, Berglund B, Niklasson B, Blennow M, Sandborgh-Englund G (2005) Transport of methylmercury and inorganic mercury to the fetus and breast-fed infant. *Environ Health Perspect* 113:1381-5

Björnberg KA, Vahter M, Petersson-Grawé K och Berglund M (2005). Methyl mercury exposure in Swedish women with high fish consumption. *Science of the total environment* 341:45-52

Björnberg KA, Vahter M, Petersson-Grawé K, Glynn A, Cnattingius S, Darnerud PO, et al (2003) Methyl mercury and inorganic mercury in Swedish pregnant women and in cord blood: influence of fish consumption. *Environmental Health Perspectives* 111:637-641

Blix-Holmberg U, Björs U, Edelsjö J, Elms N, Elinder C-G, Lind B, Lindahl L, Nilsson B, Ohlsson M, Oksztel-Nilsson B, Ottosson Å, Sundstedt K (1979) Mätning av bly i blod hos stockholmare. *Läkartidningen*. 76. 45, 3983-3984

Börjesson J, Gerhardsson L, Schütz A, Perfekt R, Mattsson S, Skerfving S (2001) Kidney cadmium as compared to other markers of cadmium exposure in workers at a secondary metal smelter. *Am J Ind Med* 39:19-28

Claeys-Thoreau F, Thiessen L, Bruaux P, Ducoffre G, Verduyn G (1987) Assessment and comparison of human exposure to lead between Belgium, Malta, Mexico and Sweden. *Int. Arch. Occup Environ Health* 59:31-41

Czub G, McLachlan MS (2004) A food chain model to predict the levels of lipophilic organic contaminants in humans. *Environ Toxicol Chem* 23:2356-66

Darnerud PO (1999) Analys av organiska miljögifter i modersmjölk. Rapport till den nationella miljöövervakningen, Naturvårdsverket, kontrakt 215 805

Darnerud PO (1999) Organiska miljögifter i modersmjölk. Naturvårdsverket rapport 5015

Darnerud PO, Atuma S, Aune M, Cnattingius S, Wernroth M-L, Wicklund-Glynn A (1998) Polybrominated diphenyl ethers (PBDEs) in breast milk from primiparous women in Uppsala county, Sweden. *Organohalogen Comp* 35:411-414

Darnerud PO, Eriksen GS, Johannesson T, Larsen PB, Viluksela M (2001) Polybrominated diphenyl ethers: occurrence, dietary exposure, and toxicology. *Environ Health Perspect* 109 Suppl 1:49-68

Darnerud PO, Wicklund Glynn A, et al (2000) Dioxiner i svensk bröstmjöl och haltsamband med PCB-kongener. Rapport till den nationella miljöövervakningen, Naturvårdsverket, kontrakt 215 807

Darnerud PO, Lignell S, Glynn A, Aune M, Törnkvist A, Stridsberg M. [POP levels in breast milk and maternal serum and thyroid hormone levels in mother-child pairs from Uppsala, Sweden.](#) *Environ Int.* 2010 Feb;36(2):180-7. Epub 2009 Dec 1. PMID: 19954849

Elinder CG, Friberg L, Lind B, Jawaid M (1983) Lead and cadmium levels in blood samples from the general population of Sweden. *Environ Res* 30:233-53

Elinder CG, Friberg L, Lind B, Nilsson B, Svartengren M, Övermark I (1985) Blyhalten i blod hos stockholmare har sjunkit mellan åren 1980 och 1984. *Läkartidningen.* 84. 49, 4298-4302

Elinder CG, Friberg L, Lind B, Nilsson B, Svartengren M, Övermark I (1986) Decreased blood lead levels in residents of Stockholm for the period 1980-1984. *Scand J Work Environ Health* 12:114-120

Elinder C-G, Kjellström (1977) Cadmium concentration in samples of human kidney cortex from the 19th century. *Ambio* 6. 5:270-272

Elinder CG, Kjellström T, Lind B, Molander ML, Silander T (1978) Cadmium concentrations in human liver, blood, and bile: comparison with a metabolic model. *Environ Res* 17:236-41

Elinder CG, Kjellström T, Linnman L, Pershagen G (1978) Urinary excretion of cadmium and zinc among persons from Sweden. *Environ Res* 15:473-84

Elinder CG, Lind B, Kjellström T, Linnman L, Friberg L (1976) Cadmium in kidney cortex, liver, and pancreas from Swedish autopsies. Estimation of biological half time in kidney cortex, considering calorie intake and smoking habits. *Arch Environ Health* 31:292-302

Elinder CG, Piscator M, Linnman L (1977) Cadmium and zinc relationships in kidney cortex, liver, and pancreas. *Environ Res* 13:432-40

Engström A, Michaëlsson K, Suwazono Y, Wolk A, Vahter M, Akesson A. Long-term cadmium exposure and the association with bone mineral density and fractures in a population-based study among women. *J Bone Miner Res*. 2010 Aug 23. [Epub ahead of print]

Eriksson S, Darnerud PO, Aune M, Bjerselius R, Slanina P, Cnattingius S och Glynn A. (2003) Syntetiska myskföreningar i bröstmjolk och fisk, Resultatrapport till Naturvårdsverket. Livsmedelsverket, Uppsala. ([http://www.imm.ki.se/Datavard/PDF/Mysk\\_mjolk.pdf](http://www.imm.ki.se/Datavard/PDF/Mysk_mjolk.pdf))

Friis L, Petersson L, Edling C (1998) Reduced cadmium levels in human kidney cortex in Sweden. *Environ. Health Perspect.* 106, 175-8

Fängström B, Athanassiadis I, Odsjö T, Norén K, Bergman Å. Temporal trends of polybrominated diphenyl ethers and hexabromocyclododecane in milk from Stockholm mothers, 1980–2004. *Mol Nutr Food Res* 2008; 52:187–93.

Gerhardsson L, Lundh T. [Metal concentrations in blood and hair in pregnant females in southern Sweden](#). *J Environ Health*. 2010 Jan-Feb;72(6):37-41.PMID: 20104833

Glynn AW, Wolk A, Aune M, Atuma S, Zettermark S, Maehle-Schmid M, Darnerud PO, Becker W, Vessby B, Adami HO (2000) Serum concentrations of organochlorines in men: a search for markers of exposure. *Sci Total Environ* 18;263:197-208

Glynn AW, Atuma S, Aune M, Darnerud PO, Cnattingius S (2001) Polychlorinated biphenyl congeners as markers of toxic equivalents of polychlorinated biphenyls, dibenzo-p-dioxins and dibenzofurans in breast milk. *Environ Res* 86:217-28

Glynn AW, Darnerud PO, Aune M, Bjerselius R, Baumann B, Cnattingius S (2003) PCBs and dioxins in breast milk - levels and trends in Sweden 1996-2001. Abstract to Dioxin 2003

Glynn AW, Granath F, Aune M, Atuma S, Darnerud PO, Bjerselius R, Vainio H, Weiderpass E (2003) Organochlorines in Swedish women: determinants of serum concentrations. *Environ Health Perspect* 111:349-55

Glynn A, Aune M, Darnerud PO, Atuma S, Cnattingius S, Bjerselius R, Becker W, Lind Y. Studie av förstföderskor. Organiska miljögifter hos gravida och ammande. Del 1 Serumnivåer. Rapport 4 – 2006. Livsmedelsverket, Uppsala.

Glynn A, Aune M, Darnerud PO, Cnattingius S, Bjerselius R, Becker W, Lignell S. Determinants of serum concentrations of organochlorine compounds in Swedish pregnant women: a cross-sectional study. *Environ Health*. 2007 Feb 1;6:2

Glynn A, Lignell S, Darnerud PO, Aune M, Halldin Ankarberg E, Bergdahl IA, Barregård L, Bensryd I. Regional differences in levels of chlorinated and brominated pollutants in mother's milk from primiparous women in Sweden. *Environ Int*. 2010 Aug 28. [Epub ahead of print]

Granvik M, Sandberg B-M, Friberg L, Lind B, Nilsson B (1988) Låga halter av bly och kadmium i blod hos skolbarn i Dalarna. *Läkartidningen*. 85. 5, 320-322

Hagmar L (1997) Konsumtion av fet östersjöfisk och PCB i blod hos svenska kvinnor, i Naturvårdsverkets rapport 4760

Hagmar L, Björk J, Sjödin A, Bergman Å, Erfurth EM (2001) Plasma levels of persistent organohalogenes and hormone levels in adult male humans. *Arch Environ Health* 56:138-43

Hagmar L, Rylander L, Dyremark E, Klasson-Wehler E, Erfurth EM (2001) Plasma concentrations of persistent organochlorines in relation to thyrotropin and thyroid hormone levels in women. *Int Arch Occup Environ Health* 74:184-8

Hallén IP, Jorhem L, Lagerkvist BJ, Oskarsson A (1995) Lead and cadmium levels in human milk and blood. *Sci Tot Environ* 166:149-155

Hanberg A, Ahlborg U (1994) PCB och tungmetaller hos kvinnor i Örebro län. IMM-rapport

Harrad S, de Wit CA, Abdallah MA, Bergh C, Björklund JA, Covaci A, Darnerud PO, de Boer J, Diamond M, Huber S, Leonards P, Mandalakis M, Ostman C, Haug LS, Thomsen C, Webster TF. [Indoor contamination with hexabromocyclododecanes, polybrominated diphenyl ethers, and perfluoroalkyl compounds: an important exposure pathway for people?](#)

*Environ Sci Technol*. 2010 May 1;44(9):3221-31. Review.

Hovander L, Malmberg T, Athanasiadou M, Athanassiadis I, Rahm S, Bergman Å, Wehler EK (2002) Identification of hydroxylated PCB metabolites and other phenolic halogenated pollutants in human blood plasma. *Arch Environ Contam Toxicol* 42:105-17

- Huang W, Åkesson B, Svensson BG, Schütz A, Burk RF, Skerfving S (1995) Selenoprotein P and glutathione peroxidase (EC 1.11.1.9) in plasma as indices of selenium status in relation to the intake of fish. *Br J Nutr* 73:455-61
- Högberg J, Hanberg A, Berglund M, Skerfving S, Remberger M, Calafat A.M, Falk Filipsson A, Jansson B, Johansson N, Appelgren M, Håkansson H. Phthalate diesters and their metabolites in human breast milk, blood and urine as biomarkers of exposure in vulnerable populations. *Environ Health Perspect* Doi: 10.1289/ehp.10788 (Online 21 December 2007).
- Jakobsson K, Athanasiadou M, Christiansson A, Athanassiadis I, Bergman Å, Hagmar L (2005) Polybrominated diphenyl ethers (PBDEs) in serum from Swedish men 1988-2002. A longitudinal study. *Yrkes- och miljömedicin, Lund. Projekt 215 04 11*
- Jarosińska D, Horvat M, Sällsten G, Mazzolai B, Dabkowska B, Prokopowicz A, Biesiada M, Barregård L. Urinary mercury and biomarkers of early renal dysfunction in environmentally and occupationally exposed adults: a three-country study. *Environ Res.* 2008 Oct;108(2):224-32.
- Jawaid M, Lind B, Elinder C-G (1983) Determination of cadmium in urine by extraction and flameless atomic-absorption spectrophotometry. *Talanta* 30:509-513
- Johnsson C, Schütz A, Sällsten G (2005) Impact of consumption of freshwater fish on mercury levels in hair, blood, urine, and alveolar air. *J Toxicol Environ Health A* 22;68:129-40
- Johnsson C, Sällsten G, Schütz A, Sjörs A, Barregård L. (2004) Hair mercury levels versus freshwater fish consumption in household members of Swedish angling societies. *Environmental Research* 96:257-263
- Järup L, Carlsson MD, Elinder CG, Hellström L, Persson B, Schütz A (1995) Enzymuria in a population living near a cadmium battery plant. *Occup Environ Med* 52:770-772
- Järup L, Hellström L, Alfven T, Carlsson MD, Grubb A, Persson B, Pettersson C, Spang G, Schütz A, Elinder CG (2000) Low level exposure to cadmium and early kidney damage: the OSCAR study. *Occup Environ Med* 57:668-72. Erratum in: *Occup Environ Med* 2002 ;59:497
- Kaj L, Andersson J, Palm Cousins A, Remberger M, Brorström-Lundén E, Cato I (2005) Results from the Swedish National Screening Programme 2004. Subreport 4: Siloxanes. *IVL-rapport B1643*
- Kjellström T (1979) Exposure and accumulation of cadmium in populations from Japan, the United States, and Sweden. *Environ Health Perspect* 28:169-197



Kärrman A, Ericson I, van Bavel B, Darnerud PO, Aune M, Glynn A, Lignell S, Lindström G. Exposure of perfluorinated chemicals through lactation: levels of matched human milk and serum and a temporal trend, 1996-2004, in Sweden. *Environ Health Perspect.* 2007 Feb;115(2):226-30.

Lagerkvist B (1996) Studie av bakgrundshalter av miljö- och hälsofarliga metaller och organiska miljögifter hos människor. Naturvårdsverket, rapport 4902

Lagerkvist BJ, Ekesrydh S, Englyst V, Nordberg GF, Söderberg HA, Wiklund DE (1996) Increased blood lead and decreased calcium levels during pregnancy: a prospective study of Swedish women living near a smelter. *Am J Publ Health* 86:1247-1252

Lagerkvist BJ, Nordberg GF, Söderberg HA, Ekesrydh S, Englyst V, Gustavsson M, Gustavsson NO, Wiklund DE (1992) Placental transfer of cadmium. *IARC Sci Publ No* 118:287-291

Lagerkvist BJ, Sandberg S, Frech W, Jin T, Nordberg GF (1996) Is placenta a good indicator of cadmium and lead exposure? *Arch Environ Health* 51:389-394

Lagerkvist BJ, Söderberg HA, Nordberg GF, Ekesrydh S, Englyst V (1993) Biological monitoring of arsenic, lead and cadmium in occupationally and environmentally exposed pregnant women. *Scand J Work Environ Health* 19 Suppl 1:50-53

Lagerkvist BJ, Söderberg HÅ, Ekesrydh S, Englyst V, Gustavssin M, Nordberg G (199?) Bly- kadmium- och arsenikhalter hos gravida i Rönnskärsområdet. Umeå Universitet

Langworth S, Elinder CG, Göthe CJ, Vesterberg O (1991) Biological monitoring of environmental and occupational exposure to mercury. *Int Arch Occup Environ Health* 63:161-7

Larsdotter M, Darnerud PO, Aune M, Glynn A, Bjerselius R (2005) Serumnivåer av pentaklorfenol, polyklorerade bifenyler och hydroxylerade metaboliter av PCB under graviditet och amning. Livsmedelsverket. Rapport till Naturvårdsverket, <http://www.naturvardsverket.se/dokument/mo/modok/export/klorfenoler.pdf>

Lignell S, Aune M, Darnerud PO, Glynn A (2004) Kompletterande mätningar av syntetiska myskföreningar i bröstmjolk från förstföderskor i Uppsala, 1996-2003. Livsmedelsverket. Rapport till Naturvårdsverket. [http://www.naturvardsverket.se/dokument/mo/modok/export/mysk\\_brostmjolk.pdf](http://www.naturvardsverket.se/dokument/mo/modok/export/mysk_brostmjolk.pdf)

Lignell S, Glynn A, Darnerud PO, Aune M, Bergdahl I, Barregård L, Bensryd I (2005) Regionala skillnader i halter av persistenta organiska miljögifter i bröstmjolk från förstföderskor i Uppsala, Göteborg, Lund och Lycksele. Livsmedelsverket. Rapport till Naturvårdsverket, <http://www.naturvardsverket.se/dokument/mo/modok/export/regional%20trend.pdf>

Lignell S, Darnerud PO, Aune M, Cnattingius S, Hajslova J, Setkova L, Glynn A. Temporal trends of synthetic musk compounds in mother's milk and associations with personal use of perfumed products. *Environ Sci Technol*. 2008 Sep 1;42(17):6743-8

Lignell S, Aune M, Darnerud PO, Cnattingius S, Glynn A. [Persistent organochlorine and organobromine compounds in mother's milk from Sweden 1996-2006: compound-specific temporal trends](#). *Environ Res*. 2009 Aug;109(6):760-7. Epub 2009 May 27. PMID: 19477439

Lind Y, Darnerud PO, Atuma S, Aune M, Becker W, Bjerselius R, Cnattingius S, Glynn A (2003) Polybrominated diphenyl ethers in breast milk from Uppsala county, Sweden. *Environ Res* 93:186-194

Lindberg A, Björnberg KA, Vahter M, Berglund M (2004) Exposure to methylmercury in non-fish-eating people in Sweden. *Environ Res* 96:28-33

Lundén A, Norén K (1998) Polychlorinated naphthalenes and other organochlorine contaminants in Swedish human milk, 1972-1992. *Arch Environ Contam Toxicol* 34:414-423

Lutz E, Lind B, Herin P, Krakau I, Bui TH, Vahter M (1996) Concentrations of mercury, cadmium and lead in brain and kidney of second trimester fetuses and infants. *J Trace Elem Med Biol* 10:61-67

Länsstyrelsen i Örebro län, miljöenheten (1995) PCB och tungmetaller hos kvinnor i Örebro län, publikation 1995:2

Meironyte D, Noren K, Bergman Å (1999) Analysis of polybrominated diphenyl ethers in Swedish human milk. A time-related trend study, 1972-1997. *J Toxicol Environ Health A* 26;58:329-41

Molin M, Schütz A, Skerfving S, Sällsten G (1991) Mobilized mercury in subjects with varying exposure to elemental mercury vapour. *Int Arch Occup Environ Health* 63:187-92

Naturvårdsverket (1995) PCB och tungmetaller hos kvinnor i Örebro län, ur Rapport 4555

Nilsson H, Kärrman A, Westberg H, Rotander A, van Bavel B, Lindström G. [A time trend study of significantly elevated perfluorocarboxylate levels in humans after using fluorinated ski wax](#).

Nilsson U, Schütz A, Skerfving S, Mattsson S (1995) Cadmium in kidneys in Swedes measured in vivo using X-ray fluorescence analysis. *Int Arch Occup Environ Health* 67:405-11

Nilsson U, Skerfving S (1993) In vivo x-ray fluorescence measurements of cadmium and lead. *Scand J Work Environ Health* 19 Suppl 1:54-8

Nordberg M, Winblad B, Basun H (2000) Cadmium concentration in blood in an elderly urban population. *Biometals* 13:311-7

Nordberg M, Winblad B, Fratiglioni L, Basun H (2000) Lead concentrations in elderly urban people related to blood pressure and mental performance: results from a population-based study. *Am J Ind Med* 38:290-4

Norén K (1993) Contemporary and retrospective investigations of human milk in the trend studies of organochlorine contaminants in Sweden. *Sci Tot Environ* 139-140:347-355

Norén K, Lundén A (1991) Trend studies of polychlorinated biphenyls, dibenzo-p-dioxins and dibenzofurans in human milk. *Chemosphere* 23:1895-1901

Norén K, Lundén A, Pettersson E, Bergman Å (1996) Methylsulfonyl metabolites of PCBs and DDE in human milk in Sweden, 1972-1992. *Environ Health Perspect* 104:766-772

Norén K, Meironyté D (2000) Certain organochlorine and organobromine contaminants in Swedish human milk in perspective of past 20-30 years. *Chemosphere* 40:1111-1123

Olsson IM (2002) Biomonitoring of cadmium in cattle, pigs and humans. Akademisk avhandling, Sveriges Lantbruksuniversitet

Olsson IM, Bensryd I, Lundh T, Ottosson H, Skerfving S, Oskarsson A (2002) Cadmium in blood and urine--impact of sex, age, dietary intake, iron status, and former smoking--association of renal effects. *Environ Health Perspect* 110:1185-90

Oskarsson A, Lagerkvist BJ, Ohlin B, Lundberg K (1994) Mercury levels in the hair of pregnant women in a polluted area in Sweden. *Sci Tot Environ* 151:29-35

Oskarsson A, Ohlin B, Lagerkvist BJ (1991) Kvicksilverhalt i hår hos gravida kvinnor i Rönnskärsområdet. Livsmedelsverket, Umeå Universitet

Oskarsson A, Ohlin B, Ohlander EM, Albanus L (1990) Mercury levels in hair from people eating large quantities of Swedish freshwater fish. *Food Add Contam* 7:555-562

Oskarsson A, Palminger Hallén I, Sundberg J (1995) Exposure to toxic elements via breast milk. *Analyst* 120:765-770

Oskarsson A, Schütz A, Skerfving S, Hallen IP, Ohlin B, Lagerkvist BJ (1996) Total and inorganic mercury in breast milk in relation to fish consumption and amalgam in lactating women. *Arch Environ Health* 51:234-241

Oskarsson A, Widell A, Olsson IM, Grawe KP (2004) Cadmium in food chain and health effects in sensitive population groups. *Biometals* 17:531-4

Osman K, Åkesson A, Berglund M, Bremme K, Schütz A, Ask K, Vahter M (2000) Toxic and essential elements in placentas of Swedish women. *Clin Biochem* 33:131-138

Rappe C (1994) Sources of and human exposure to PCDDs and PCDFs. In: Banbury report 35: Biological basis for risk assessment of dioxins and related compounds, eds Gallo, Scheuplein and van der Heijden Cold Spring Harbor Laboratory Press, New York 121-129

Remberger M, Andersson J, Palm Cousins A, Kaj L, Ekheden Y, Dusan B, Brorström-Lundén E, Cato I (2005) Results from the Swedish National Screening Programme 2004. Subreport 1: Adipates. IVL-rapport B1645

Rignell-Hydbom A, Rylander L, Giwercman A, Jonsson BA, Lindh C, Eleuteri P, Rescia M, Leter G, Cordelli E, Spano M, Hagmar L (2005) Exposure to PCBs and p,p'-DDE and human sperm chromatin integrity. *Environ Health Perspect* 113:175-9

Rignell-Hydbom A, Axmon A, Lundh T, Jönsson BA, Tiido T, Spano M. Dietary exposure to methyl mercury and PCB and the associations with semen parameters among Swedish fishermen. *Environ Health*. 2007 May 8;6:14.

Rignell-Hydbom A, Skerfving S, Lundh T, Lindh CH, Elmståhl S, Bjellerup P, Jönsson BA, Strömberg U, Åkesson A. [Exposure to cadmium and persistent organochlorine pollutants and its association with bone mineral density and markers of bone metabolism on postmenopausal women.](#) *Environ Res*. 2009 Nov;109(8):991-6. Epub 2009 Sep 5. PMID: 19733845

Rylander L, Dyremark E, Strömberg U, Östman C, Hagmar L (1997) The impact of age, lactation and dietary habits on PCB in plasma in Swedish women. *Sci Total Environ* 207:55-61

Rylander L, Strömberg U, Dyremark E, Östman C, Nilsson-Ehle P, Hagmar L (1998) Polychlorinated biphenyls in blood plasma among Swedish female fish consumers in relation to low birth weight. *Am J Epidemiol* 147:493-502

Rylander L, Hagmar L, Wallin E, Sjöström AK, Tysklind M. [Intra-individual variations and temporal trends in dioxin levels in human blood 1987-2002.](#) *Chemosphere*. 2009 Sep;76(11):1557-62. Epub 2009 Jun 25. PMID: 19559463

Sandborgh Englund G, Adolfsson-Erici M, Ekstrand J, Odham G. Triclosan: Biotillgänglighet och farmakokinetik. Resultatrapport, kontrakt 219 0206, Naturvårdsverket

Schütz A, Attewell R, Skerfving S (1989) Decreasing blood lead in Swedish children, 1978-1988. *Arch Environ Health* 44:391-394

Schütz A, Bergdahl IA, Ekholm A, Skerfving S (1996) Measurement by ICP-MS of lead in plasma and whole blood of lead workers and controls. *Occup Environ Med* 53:736-40

Seldén, A., Bergström, B., Ek, B.-M., Öjstrand, B.-M., Edlund, B., Högdahl, C., Norberg, C., Ohlson, C.-G., Lundholm, C., Weiss, L. & Andersson, L., 2006. Uran i dricksvatten från bergborrade brunnar – exponering, utsöndring och njurtoxiska effekter. Rapport från Yrkesmedicin YM 5/06. Yrkes- och miljömedicinska kliniken. Universitetssjukhuset Örebro.

Seldén AI, Lundholm C, Edlund B, Högdahl C, Ek BM, Bergström BE, Ohlson CG. Nephrotoxicity of uranium in drinking water from private drilled wells. *Environ Res* 2009;109:486-94.

Sjödin A, Hagmar L, Klasson-Wehler E, Björk J, Bergman Å (2000) Influence of the consumption of fatty Baltic Sea fish on plasma levels of halogenated environmental contaminants in Latvian and Swedish men. *Environ Health Perspect* 108:1035-41  
Sjödin A, Hagmar L, Klasson-Wehler E, Kronholm-Diab K, Jakobsson E, Bergman Å (1999) Flame retardant exposure: polybrominated diphenyl ethers in blood from Swedish workers. *Environ Health Perspect* 107:643-648

Sjödin A, Patterson DG Jr, Bergman Å (2003) A review on human exposure to brominated flame retardants--particularly polybrominated diphenyl ethers. *Environ Int* 29:829-39

Skerfving S (1988) Blyhalt i blod hos sydsvenska barn 1978-87. Slutrapport för Statens Naturvårdsverks projekt 611-644-1985-UF.

Skerfving S (1988) Mercury in women exposed to methylmercury through fish consumption, and in their newborn babies and breast milk. *Bull Environ Contam Toxicol* 41:475-482.

Skerfving S, Nilsson U (1992) Assessment of accumulated body burden of metals. *Toxicol Lett* 64-65 Spec No 17-24

Skerfving S, Nilsson U, Schutz A, Gerhardsson L (1993) Biological monitoring of inorganic lead. *Scand J Work Environ Health* 19 Suppl 1:59-64.

Skerfving S, Schütz A, Ranstam J (1986) Decreasing lead exposure in Swedish children, 1978-84. *Sci Total Environ* 58:225-9

Skerfving S, Strömberg U, Schütz A (1997) Blyexponeringens långsiktiga trend inom den nationella miljöövervakningen. Uppdatering 1997. Rapport för Naturvårdsverkets projekt 215 507B

Skerfving S, Svensson BG, Asplund L, Hagmar L (1994) Exposure to mixtures and congeners of polychlorinated biphenyls. Clin Chem 40:1409-15

Smolders R, Alimonti A, Cerna M, Den Hond E, Kristiansen J, Palkovicova L, Ranft U, Seldén AI, Telisman S, Schoeters G. [Availability and comparability of human biomonitoring data across Europe: a case-study on blood-lead levels.](#) Sci Total Environ. 2010 Feb 15;408(6):1437-45.

Stroh E, Lundh T, Oudin A, Skerfving S, Strömberg U. [Geographical patterns in blood lead in relation to industrial emissions and traffic in Swedish children, 1978-2007.](#) BMC Public Health. 2009 Jul 10;9:225.PMID: 19591669

Strömberg U, Schütz A, Skerfving S (1995) Substantial decrease of blood lead in Swedish children, 1978-94, associated with petrol lead. Occup Environ Med 52:764-769

Strömberg U, Lundh T, Skerfving S (2002) Låga blyhalter hos barn. Bulletin från Yrkes- och miljömedicin i Lund 20(4), 13

Strömberg U, Lundh T, Skerfving S. Yearly measurements of blood lead in Swedish children since 1978: the declining trend continues in the petrol-lead-free period 1995-2007 Environ Res. 2008 Jul;107(3):332-5.

Sundström M, Ehresman DJ, Bignert A, Butenhoff JL, Olsen GW, Chang SC, Bergman A. [A temporal trend study \(1972-2008\) of perfluorooctanesulfonate, perfluorohexanesulfonate, and perfluorooctanoate in pooled human milk samples from Stockholm, Sweden.](#) Environ Int. 2010 Sep 27. [Epub ahead of print] PMID: 20880590

Svartengren M, Elinder CG, Friberg L, Lind B (1986) Distribution and concentration of cadmium in human kidney. Environ Res 39:1-7

Svartengren M, Friberg L, Lind B (1991) Assessment of human exposure to lead. Comparison of exposure between Belgium, Malta, Mexico and Sweden. HK Dillon, MH Ho (eds) Biological Monitoring of Exposure to Chemicals: Metals XIX, John Wiley and sons, New York, 73-84

Svensson BG, Nilsson A, Hansson M, Rappe C, Åkesson B, Skerfving S (1991) Exposure to dioxins and dibenzofurans through the consumption of fish. New England J Med 324: 8-12

Svensson BG, Schütz A, Nilsson A, Skerfving S (1992) Lead exposure in indoor firing ranges. Int Arch Occup Environ Health 64:219-221

Svensson BG, Schütz A, Nilsson A, Åkesson I, Åkesson B, Skerfving S (1992) Fish as a source of exposure to mercury and selenium. Sci Tot Environ 126:61-74

Sällsten G, Barregård L (1997) Uppföljning av kvicksilver i blod hos gravida kvinnor i Göteborg. Länsstyrelsen i V Götalands län

Törneman N, Thuresson C. Screening of platinum group metals; Pt, Rh and Pl. Screening Report 2007:2. Rapport till Naturvårdsverket. SWECO VIAK, Malmö 2007

Törneman N, Thuresson C. Screening of platinum group metals; Pt, Rh and Pl. Screening Report 2007:2. Appendix 1, sampling localities in the Stockholm area. Rapport till Naturvårdsverket. SWECO VIAK, Malmö 2007

Törneman N (2010). Exposure and effect screening in urine of women: 1. Metals and metabolites of phthalates, organophosphate pesticides and PAHs. 2. Endocrine disturbing effects. SWECO Environment AB, Malmö.

Ulander A och Axelsson O (1974) Measurement of blood-cadmium levels. Lancet, April 13, 682-683. (Ingår i Vahter 1982)

Vahter M (1982) Assessment of human exposure to lead and cadmium through biological monitoring. National Swedish Institute of Environmental Medicine and Department of Hygiene, Karolinska Institutet, Stockholm

Vahter M, Berglund M, Friberg L, Jorhem L, Lind B, Slorach S, Åkesson A (1990) Dietary intake of lead and cadmium in Sweden. Vår Föda 42. Suppl. 2

Vahter M, Berglund M, Lind B, Jorhem L, Slorach S, Friberg L (1991) Personal monitoring of lead and cadmium exposure-a Swedish study with special reference to methodological aspects. Scand J Work Environ Health 17:65-74

Vahter M, Berglund M, Nermell B, Åkesson A (1996) Bioavailability of cadmium from shellfish and mixed diet in women. Toxicol App Pharmacol 136:332-41

Vahter M, Berglund M, Slorach S, Friberg L, Saric M, Zheng XQ, Fujita M (1991) Methods for integrated exposure monitoring of lead and cadmium. Environ Res 56:78-89

Vahter M, Berglund M, Slorach S, Jorhem L, Lind B (1992) Integrated personal monitoring of cadmium exposure in Sweden. IARC Sci Publ No 118:113-119

Vahter M, Lind B (1986) Concentrations of arsenic in urine of the general population in Sweden. Sci Tot Environ 54:1-12

Vahter M, Åkesson A, Lind B, Björs U, Schütz A, Berglund M (2000) Longitudinal study of methylmercury and inorganic mercury in blood and urine of pregnant and lactating women, as well as in umbilical cord blood. Environ Res 84:186-94

Van Oostdam JC, Dewailly E, Gilman A, Hansen JC, Odland JO, Chashchin V, Berner J, Butler-Walker J, Lagerkvist BJ, Olafsdottir K, Soininen L, Bjerregard P, Klopov V, Weber JP (2004) Circumpolar maternal blood contaminant survey, 1994-1997 organochlorine compounds. *Sci Total Environ* 1;330:55-70

Vaz R (1995) Average Swedish dietary intakes of organochlorine contaminants via food of animal origin and their relation to levels in human milk, 1975-90. *Food Add Contam* 12:542-558

Weistrand C, Norén K (1998) Polychlorinated naphthalenes and other organochlorine contaminants in human adipose and liver tissue. *J Toxicol Environ Health* 53:293-311

Willers S, Axmon A, Feyerabend C, Nielsen J, Skarping G, Skerfving S (2000) Assessment of environmental tobacco smoke exposure in children with asthmatic symptoms by questionnaire and cotinine concentrations in plasma, saliva, and urine. *J Clin Epidemiol* 53:715-21

Willers S, Schütz A, Attewell R, Skerfving S (1988) Relation between lead and cadmium in blood and the involuntary smoking of children. *Scand J Work Environ Health* 14:385-389

Zetterlund B, Winberg J, Lundgren G, Johansson G (1977) Lead in umbilical cord blood correlated with the blood lead of mother in areas with low, medium or high atmospheric pollution. *Acta Paediatr Scand* 66:167-175

Åkesson A (2000) Cadmium exposure and iron status. Akademisk avhandling. Karolinska Institutet

Åkesson A, Berglund M, Schütz A, Bjellerup P, Bremme K, Vahter M (2002) Cadmium exposure in pregnancy and lactation in relation to iron status. *Am J Public Health* 92:284-287

Åkesson A, Lundh T, Vahter M, Bjellerup P, Lidfeldt J, Nerbrand C, et al. Tubular and glomerular kidney effects in Swedish women with low environmental cadmium exposure. *Environ Health Perspect* 2005;113(11):1627-31.

Åkesson A, Julin B, Wolk A. Dietary cadmium exposure and postmenopausal endometrial cancer incidence: A population-based prospective cohort study. *Cancer Res*, 2008;68:6435-41.